

LA-UR-21-28078

Approved for public release; distribution is unlimited.

Title: Neural Density Estimation and Uncertainty Quantification for ChemCam Spectra

Author(s): Kontolati, Katiana
Panda, Nishant
Klein, Natalie Elizabeth
Moore, Juston Shane
Oyen, Diane Adele

Intended for: Report

Issued: 2021-08-12

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Neural Density Estimation and Uncertainty Quantification for ChemCam Spectra

AML Fellow: Katiana Kontolati
Ph.D. student, Johns Hopkins University

Mentors: Nishant Panda, Natalie Klein, Juston Moore, Diane Oyen

Outline

- **Background:** ChemCam data and literature review
- **Motivation:** Uncertainty quantification for ChemCam
- **Methods:** Generative modeling via normalizing flows
- **Proposed approach and Results**
- **Discussion and Conclusions**

Background: ChemCam






Illustration: ChemCam firing laser







- The ChemCam instrument of Curiosity uses laser-induced breakdown spectroscopy (LIBS)
- Fires a laser at target, vaporizes rock surfaces, creating a plasma
- Three spectrographs divide the plasma light into wavelengths for chemical analysis
- The three wavelength ranges: Ultraviolet, Violet, Visible Near-Infrared

Background

- Regression methods (SVR, PCR, CNN) have been employed for calibration (prediction of the elemental composition of samples)
- However, labeled ChemCam samples are limited


Motivation

- Focus on unsupervised learning and employ *generative models* from ChemCam analysis
- Use labels (supervised) in combination to the generative model to compute uncertainties related to predictions



Comparison of 10 regression models

Boucher, T. F., et al., (2015). *Spectr. Acta Part B: Atomic Spectroscopy*



Learning the chemical content of samples (regression results)

Castorena, J. et al., (2021). *Spectr. Acta Part B: Atomic Spectroscopy*

uncertainty propagation



Outline

- **Background:** ChemCam data and literature review
- **Motivation:** Uncertainty quantification for ChemCam
- **Methods:** Generative modeling via normalizing flows
- **Proposed approach and Results**
- **Discussion and Conclusions**

Methods


Generative modeling

- In generative modeling, any kind of observed dataset \mathcal{D} , is a finite set of samples generated from an *underlying distribution*
- The goal of any generative model is to approximate this data distribution given access to the dataset \mathcal{D}
- If we are able to *learn* a good generative model, we can use the learned model for downstream *inference*
- Perform: Sampling, density estimation, detect outliers, fill in incomplete data, representation learning




Methods: Normalizing flows (NF)

Definition: A Normalizing Flow is a transformation of a simple probability distribution (e.g., a standard normal) into a more complex distribution by a sequence of invertible and differentiable mappings.



- Mappings $f_i(z_{i-1})$ need to be computationally efficient but also expressive enough
- NF produce tractable distributions where both sampling and density evaluation can be efficient and exact
- Compared to other approaches (VAE, GAN) NF allows for exact evaluation of densities and efficient sampling
- Parallel WaveNet¹ model is currently used by Google Assistant to generate realistic speech


Transformation example



4-step flow transforming samples from a standard-normal base density to a cross-shaped target density (Papamakarios et. al., 2021 arXiv 1912.02762)

¹Oord, A., et al. (2018) *ICML* (pp. 3918-3926)

Methods: Normalizing flows (NF)



- Let us consider a directed, latent-variable model over observed variables X and latent variables Z
- In a **normalizing flow model**, the mapping between Z and X , given by $f_{\theta}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is deterministic and invertible such that $X = f_{\theta}(Z)$ and $Z = f_{\theta}^{-1}(X)$
- Using change of variables, the marginal likelihood $p(\mathbf{x})$ is given by

$$p_X(\mathbf{x}; \theta) = p_Z(f_{\theta}^{-1}(\mathbf{x})) \left| \det \left(\frac{\partial f_{\theta}^{-1}(\mathbf{x})}{\partial \mathbf{x}} \right) \right|$$

- “**Normalizing**”: change of variables gives a normalized density after applying an invertible transformation.
- “**Flow**”: invertible transformations can be composed to create more complex invertible transformations.



Outline

- **Background:** ChemCam data and literature review
- **Motivation:** Uncertainty quantification for ChemCam
- **Methods:** Generative modeling via normalizing flows
- **Proposed approach and Results**
- **Discussion and Conclusions**


Dimension reduction

Non-negative matrix factorization (NMF)


- Assume random vector $y \in \mathbb{R}^M$ where $Y = [y_1, \dots, y_N] \in \mathbb{R}^{N \times M}$
- $M = 5606$ (original dimensionality)
- Decompose $Y \approx XV$
- $X \in \mathbb{R}_{\geq 0}^{N \times L}$ and $V \in \mathbb{R}_{\geq 0}^{L \times M}$, $L \ll M$
- X : non-negative basis matrix, V : non-negative coefficient matrix
- Optimization: Minimize the Frobenius norm between Y and XV



original space
 $y \in \mathbb{R}^M$



NMF
latent space
 $x \in \mathbb{R}^L$






original space
 $y \in \mathbb{R}^M$

Normalizing flow model

\mathbf{x} : Spectral latent variable
 \mathbf{z} : Latent variable (\sim normal)

Latent space dimension: 15



Train normalizing flow model on the latent space

Real-NVP (Real-valued non-volume preserving)

Forward flow: $\mathbf{x}_{1:d} = \mathbf{z}_{1:d}$


$$\mathbf{x}_{d+1:D} = \mathbf{z}_{d+1:D} \odot \exp(f_\alpha(\mathbf{z}_{1:d})) + f_\mu(\mathbf{z}_{1:d}),$$

Inverse flow: $\mathbf{z}_{1:d} = \mathbf{x}_{1:d}$

$$\mathbf{z}_{d+1:D} = (\mathbf{x}_{d+1:D} - f_\mu(\mathbf{x}_{1:d}) \odot \exp(-f_\alpha(\mathbf{x}_{1:d})))$$

Determinant of Jacobian:

$$\det(J) = \prod_{i=1}^{D-d} \exp(f_\alpha(\mathbf{z}_{1:d}))_i = \exp\left(\sum_{i=1}^{D-d} f_\alpha(\mathbf{z}_{1:d})_i\right)$$






Single coupling flow architecture

Dinh et al. (2016) Density estimation using real nvp. arXiv:1605.08803

Normalizing flow model

Latent space dimension: 15



Train normalizing flow model

x : Spectral latent variable
 z : Latent variable (\sim normal)

Real-NVP

Dinh et al. (2016) Density estimation using real nvp. arXiv:1605.08803

Forward flow:


$$\mathbf{x}_{1:d} = \mathbf{z}_{1:d}$$

$$\mathbf{x}_{d+1:D} = \mathbf{z}_{d+1:D} \odot \exp(f_\alpha(\mathbf{z}_{1:d})) + f_\mu(\mathbf{z}_{1:d}),$$

Inverse flow:





$$\mathbf{z}_{1:d} = \mathbf{x}_{1:d}$$

$$\mathbf{z}_{d+1:D} = (\mathbf{x}_{d+1:D} - f_\mu(\mathbf{x}_{1:d}) \odot \exp(-f_\alpha(\mathbf{x}_{1:d})))$$



$$z \sim N(0, I_{15})$$


generative direction
 normalizing direction



$$x \sim p_x(x)$$


Map LIBS spectra to compositions

Train a MLP model for each oxide



w.t.% oxides

Target	SiO2	TiO2	Al2O3	FeOT	MgO	CaO	Na2O	K2O
AGV2	59.30	1.05	16.91	6.019720	1.79	5.200	4.19	2.88
BCR-2	54.10	2.26	13.50	12.417359	3.59	7.120	3.16	1.79
BEN	38.20	2.61	10.07	11.610990	13.15	13.870	3.18	1.39
BHVO2	49.90	2.73	13.50	11.067646	7.23	11.400	2.22	0.52
BIR-1a	47.70	0.97	15.40	10.193606	9.70	13.400	1.81	0.03
...











Accuracy of results

$$R^2 = 1 - \frac{RSS}{TSS}$$

RSS: sum of squares of residuals


TSS: total sum of squares



Uncertainty quantification via bootstrapping

Bootstrapping: Statistics resampling method that assigns measures of accuracy for any sample estimate

For a new sample $\mathbf{y}_0 \in \mathbb{R}^M$ we get a prediction: $\mathbf{v}_0 = \underbrace{v^{(i)}(\mathbf{y}_0)}_{\text{model uncertainty}} + \underbrace{r(\mathbf{y}_0)}_{\text{data uncertainty}}$



Predictions with uncertainty for 15 random samples

Uncertainty quantification via bootstrapping

Evaluation of prediction intervals

Coverage: The rate at which the *actual* values fall within the range of the prediction interval


Table. Coverage results (95% confidence intervals)

oxide	# of test samples	# of covered samples	Coverage (%)
SiO ₂	139	118	86.33
TiO ₂	139	137	98.56
Al ₂ O ₃	139	120	86.33
FeO _T	139	120	86.33
MgO	139	120	86.33
CaO	139	134	96.40
Na ₂ O	139	130	93.53
K ₂ O	139	125	89.93

Predictions with uncertainty for novel samples



- Samples generated by the normalizing flow model
- Real samples collected on Mars from ChemCam

Data generation





or...

ChemCam



Prediction with uncertainty



Outline

- **Background:** ChemCam data and literature review
- **Motivation:** Uncertainty quantification for ChemCam
- **Methods:** Generative modeling via normalizing flows
- **Proposed approach and Results**
- **Discussion and Conclusions**

Conclusions

- Generative modeling can be successfully applied to model real-world data
- Normalizing flow models can be efficiently constructed on latent spaces for fast downstream inference
- Unsupervised and supervised learning can be combined to form an uncertainty quantification framework

References

1. Wiens et al. "The ChemCam instrument suite on the Mars Science Laboratory (MSL) rover: Body unit and combined system tests." *Space science reviews* 170, no. 1 (2012): 167-227.
2. Forni et al. "Independent component analysis classification of laser induced breakdown spectroscopy spectra." *Spectrochimica Acta Part B: Atomic Spectroscopy* 86 (2013): 31-41.
3. Castorena et al. "Deep spectral CNN for laser induced breakdown spectroscopy." *Spectrochimica Acta Part B: Atomic Spectroscopy* 178 (2021): 106125.
4. Kobyzev et al. "Normalizing flows: An introduction and review of current methods." *IEEE Transactions on Pattern Analysis and Machine Intelligence* (2020).
5. Kumar et al. "Bootstrap prediction intervals in non-parametric regression with applications to anomaly detection." In *Proc. 18th ACM SIGKDD Conf. Knowl. Discovery Data Mining*. 2012.
6. Papamakarios et al. "Normalizing flows for probabilistic modeling and inference." *arXiv preprint arXiv:1912.02762* (2019).
7. Dinh et al. "Density estimation using real nvp." *arXiv preprint arXiv:1605.08803* (2016).

Thank you! 